


THE IUD

99% EFFECTIVE



HOW DOES THE IUD WORK?

There are two types of intrauterine devices (IUDs). One type contains a hormone (progestin) like one your body makes. This hormone stops your ovaries from releasing eggs, makes the mucus in your cervix too thick for sperm to pass through, and thins the lining inside the uterus. The other type of IUD contains copper. Copper prevents the sperm from fertilizing the egg.

HOW DO I START THE IUD?

The IUD is inserted inside your uterus by a healthcare provider. The IUD can be inserted at any time if you are not pregnant. The copper IUD starts working right away. With hormonal IUDs, you should use condoms as a backup method for the first 7 days after it is placed. If you have unprotected sex in this first week, take emergency contraception (EC).

HOW DO I USE THE IUD?

Once it is placed, you don't need to do anything. Depending on the IUD, it will provide between 3 to 12 years of birth control. If you like it, you can get another one right away.

WHAT CAN I EXPECT?

You will feel about the same.

The following are common side effects: Many people experience cramps when the IUD is inserted. If pain with insertion is a concern, you may ask your provider for options to help you with this. You may have extra menstrual bleeding for the first few months. After this, you will feel about the same and you will not feel the IUD inside you.

With the copper IUD, your periods may become heavier or may stay about the same. With the hormonal IUD your periods may become lighter or less frequent over time, and you may have no period at all. This is normal.

WHAT DO I NEED TO DO ONCE I HAVE THE IUD?

Nothing. You should see your healthcare provider for your annual exam and sexually transmitted infection (STI) testing at anytime you would like the testing.

DOES THE IUD HAVE BENEFITS?

The IUD is safe and effective birth control.

It keeps you from getting pregnant for 3 to 12 years. It is safe to use while breastfeeding. With hormonal IUDs, periods can become lighter, less painful, or not happen at all. The IUD has no effect on your ability to get pregnant in the future. You don't have to remember to use this every day.

DOES THE IUD HAVE RISKS?

The IUD is safer than pregnancy. Serious problems are rare and usually happen in the first months. The IUD can be pushed through the wall of the uterus, you can develop an infection, or the IUD can slip out. Seek medical attention if you develop the following: heavy bleeding, cramping, pain, abnormal vaginal discharge, fever, or think you might be pregnant.

This method does not protect against sexually transmitted infections (STIs) or HIV. Always use condoms to protect yourself.


WHAT IF I HAD UNPROTECTED SEX?

With the IUD, you are protected against pregnancy. If you are thinking about an IUD and had unprotected sex in the last 5 days, you can use a copper IUD or specific hormonal IUDs for both emergency contraception (EC) and long term birth control.

You can also take EC. EC works better the sooner you take it.

THE GEL

86% EFFECTIVE



HOW DOES THE GEL WORK?

The gel is a hormone-free method of birth control that must be inserted each time you have vaginal sex. It works by lowering the pH of the vagina and decreasing the motility or mobility of the sperm. It is not a spermicide.

HOW DO I START THE GEL?

The gel is an on-demand method of birth control. It may be used during any time of your menstrual cycle. It is only to be inserted before vaginal sex. Also, you may use the gel as soon as it is safe to resume intercourse following childbirth, abortion, or miscarriage.

HOW DO I USE THE GEL?

The gel is designed to stay inside the vagina during sex. Each applicator only works for one session of sex.

First, insert the plunger into the applicator, pushing down until they connect. If gel enters the pink cap, use a new applicator. Next, remove the pink cap. The gel is now ready to use!

Bend or spread your knees apart. Gently guide the applicator into the vagina, pushing it as far back as you comfortably can. With your index finger, push the plunger until it stops. Remove the applicator from the vagina; the gel will remain inside the vagina.

Each applicator only works for one session of sex. The gel starts working immediately. A new applicator will need to be used before each session of sex. Remember: *1 dose, 1 hour, 1 act.*

WHAT CAN I EXPECT?

You will feel about the same.

The gel works locally in your vagina and is not absorbed into the rest of your body. The most common side effects are: vaginal burning and itching, yeast infection, urinary tract infection, vaginal area discomfort, and vaginal discharge.

DOES THE GEL HAVE BENEFITS?

The gel can be used with other forms of birth control such as the pill or IUDs. It should not be used with the ring. The gel has no effect on your ability to get pregnant in the future. It is safe to use while breastfeeding. You don't have to remember to use it every day if you are not having sex.

DOES THE GEL HAVE RISKS?

The gel is safer than pregnancy. Serious problems are rare. Seek medical attention if you develop the following: vaginal itching, burning while urinating, or think you may be pregnant.

This method does not protect against sexually transmitted infections (STIs) or HIV. Always use condoms to protect yourself.

WHAT IF I HAD UNPROTECTED SEX?

If you don't use the gel before sex, you could become pregnant.

If it has been less than 5 days after you had unprotected sex, take emergency contraception (EC). EC works better the sooner you take it.

THE IMPLANT

99% EFFECTIVE

HOW DOES THE IMPLANT WORK?

The birth control implant contains a hormone (progestin) like one your body makes. This hormone stops your ovaries from releasing eggs, makes the mucus in your cervix too thick for sperm to pass through, and thins the lining inside the uterus.

HOW DO I START THE IMPLANT?

The implant is inserted under the skin of your upper arm by a healthcare provider. The implant can be inserted at any time if you are not pregnant.

If it is inserted up to 5 days after the start of your period, you are protected against pregnancy right away.

If it is inserted more than 5 days after the start of your period, you should use condoms as a back-up method for the first 7 days. If you have unprotected sex in this first week, take emergency contraception (EC).

HOW DO I USE THE IMPLANT?

Once it is inserted, you don't need to do anything. It lasts up to 5 years. If you like it, you can get another one.

WHAT CAN I EXPECT?

You will feel about the same.

The following are common side effects: Many people have irregular bleeding for the first 6 to 12 months. This is unpredictable—some have no bleeding; some have regular periods; some have frequent spotting or bleeding. You can have some pain or swelling when the implant is inserted that is temporary. You will feel the implant if you touch your arm, but other people generally can't see it.

WHAT DO I NEED TO DO ONCE I HAVE THE IMPLANT

Nothing. You should see your healthcare provider for your annual exam and sexually transmitted infection (STI) testing at any time you would like the testing.

DOES THE IMPLANT HAVE BENEFITS?

The implant is safe and effective birth control.

It keeps you from getting pregnant for 5 years. It is safe to use while breastfeeding. Periods may be lighter, less painful, or not happen at all. The implant has no effect on your ability to get pregnant in the future. You don't have to remember to use it every day.

DOES THE IMPLANT HAVE RISKS?

The implant is safer than pregnancy. Serious problems are rare. Some people experience mood changes, headaches, and/or acne.

Some people have pain, bruising, or infection at the insertion site. Seek medical attention if you develop the following: redness, drainage, or warmth from the insertion site, can't feel the implant, or think you might be pregnant.

This method does not protect against sexually transmitted infections (STIs) or HIV. Always use condoms to protect yourself.

WHAT IF I HAD UNPROTECTED SEX?

With the implant, you are protected against pregnancy. If you are thinking about an implant and had unprotected sex in the last 5 days, take EC. EC works better the sooner you take it.

EMERGENCY CONTRACEPTION

(EC)



HOW DOES EC WORK?

Emergency contraception (EC) is birth control that prevents pregnancy after sex. It makes it much less likely that you will become pregnant, but it is not as good as birth control that is used before or during sex.

There are four products available for EC: the copper intrauterine device (IUD), certain hormonal IUDs, levonorgestrel pill, and ulipristal acetate pill.

The levonorgestrel EC pill contains a hormone (progestin) like one your body makes. This hormone stops your ovaries from releasing eggs. Levonorgestrel EC pills are available at a pharmacy without a prescription for everyone. They may not work as well if you weigh over 200 pounds.

The ulipristal acetate EC pill is available by prescription and works well for all people, including those that weigh over 200 pounds.

HOW DO I USE EC?

Use EC as soon as you can after unprotected sex or if a condom breaks. EC works better the sooner it is taken, but can be taken up to 5 days after unprotected sex.

For EC pills, follow the directions on the packet. Most are a single dose you take as soon as possible.

If you use ulipristal acetate EC, you will need to wait 5 days before you re-start hormonal birth control.

WHAT CAN I EXPECT?

You will feel about the same.

EC has no long-term or severe side effects and has no effect on your ability to get pregnant in the future. Some people experience nausea, vomiting, headache, dizziness, breast pain, or stomach pain the day or two after taking the pill.

EC does not cause you to bleed, but your next period may come a few days early or a few days late or you may notice some bleeding before your next period.

If your period is more than 1 week late, you should take a pregnancy test.

DOES EC HAVE BENEFITS?

EC is safe and effective. Levonorgestrel EC pills can be used by anyone. It is safe to use while breastfeeding.

DOES EC HAVE RISKS?

EC pills are safe and lower your chances of getting pregnant after unprotected sex. There are no long-term or severe side effects.

EC does not protect against sexually transmitted infections (STIs) or HIV. Always use condoms to protect yourself.

WHERE CAN I GET EC?

Levonorgestrel EC is available at pharmacies without a prescription. You can get levonorgestrel EC no matter how old you are.

Ulipristal acetate EC requires a prescription.

The copper IUD and hormonal IUD require a visit with a healthcare provider.

No matter what form of birth control you are using, it is always a good idea to keep EC on hand in case you need it.



THE SHOT

94% EFFECTIVE

HOW DOES THE SHOT WORK?

The birth control shot contains a hormone (progestin) like one your body makes. This hormone stops your ovaries from releasing eggs, makes the mucus in your cervix too thick for sperm to pass through, and thins the lining inside the uterus.

HOW DO I START THE SHOT?

There are 2 ways to start the shot. We recommend the Quick Start Method.

Quick Start: Get your shot today. Use condoms as a backup method for the first 7 days after your first shot. If you have taken ulipristal acetate emergency contraception (EC), wait 5 days before starting the shot.

Next Period: Get the shot within 7 days of the start of your next period.

HOW DO I USE THE SHOT?

You get the shot in the arm, buttocks, or stomach every three months. It can be given in the muscle or under the skin.

WHAT IF I'M LATE FOR MY NEXT SHOT?

The shot works best if you get a new shot every 11-13 weeks. If your last shot was more than 15 weeks ago, use condoms for the next 7 days as a back-up method.

WHAT CAN I EXPECT?

You will feel about the same.

The following are common side effects: Many people have bleeding between periods. After the first 2-3 shots, you may have no period at all.

Some people may feel more hungry and may gain weight while on the shot. Some people report a change in acne, mood, or headaches. These can be better or worse.

WHAT CAN I EXPECT? (continued)

Long-term use of the shot may lead to lower bone density, which improves after stopping the shot.

If you do not get enough calcium in your diet, some providers will recommend calcium/vitamin D supplements.

It is not recommended for young people to use the shot for more than 2 years without an assessment by a healthcare provider.

DOES THE SHOT HAVE BENEFITS?

The shot is safe and effective birth control.

It keeps you from getting pregnant for 3 months. It is safe to use while breastfeeding. It lowers your risk of cancer of the uterus. Periods may be lighter, less painful, or not happen at all. You don't have to remember to use it every day.

After stopping the shot, it may take some people longer to get pregnant. However, it is also possible to get pregnant right away.

DOES THE SHOT HAVE RISKS?

The shot is safer than pregnancy. Serious problems are rare. Seek medical attention if you develop the following: very heavy bleeding, or think you might be pregnant.

This method does not protect against sexually transmitted infections (STIs) or HIV. Always use condoms to protect yourself.


WHAT IF I HAD UNPROTECTED SEX?

If your last shot was more than 15 weeks ago, you could become pregnant.

If it has been less than 5 days after you have had unprotected sex, take emergency contraception (EC). EC works better the sooner you take it.

THE PILL

91% EFFECTIVE



HOW DOES THE PILL WORK?

The combined birth control pill contains hormones (estrogen and progestin) like the ones your body makes. These hormones stop your ovaries from releasing eggs, make the mucus in your cervix too thick for sperm to pass through, and thin the lining inside the uterus.

HOW DO I START THE PILL?

There are 2 ways to start the pill. We recommend the Quick Start Method.

Quick Start: Take the pill as soon as you get the pack. If you take the pill within 5 days after the start of your period, you are protected against pregnancy right away. If you take the first pill more than 5 days after the start of your period, you should use condoms as a back-up method for the first 7 days. If you have taken ulipristal acetate emergency contraception (EC), wait 5 days before starting the pill.

Next Period: Take the pill within 5 days of the start after your next period.

HOW DO I USE THE PILL?

For 28 day pill packs: take 1 pill each day at the same time. After you finish a pack of pills, immediately start a new pack the next day.

For other size pill packs: Talk to your pharmacist or healthcare provider.

WHAT IF I MISS A PILL?

I forgot *ONE* pill: Take a pill as soon as you can.

I forgot *TWO* or more pills: Take one pill as soon as you can. Take your next pill at the usual time. Use condoms for 7 days as a back-up method.

WHAT CAN I EXPECT?

You will feel about the same. The following are common side effects: In the first few months you may have nausea, bleeding between periods, and/or breast discomfort. These problems generally go away after 2-3 months. Some people are concerned about weight gain. This is uncommon with the pill. Some people report a change in their mood or headaches. These can be better or worse

DOES THE PILL HAVE BENEFITS?

The pill is safe and effective birth control.

Your periods may be regular, lighter, less painful, and shorter. You may have less acne. The pill lowers your risk of getting cancer of the uterus and ovaries. The pill has no effect on your ability to get pregnant in the future.


DOES THE PILL HAVE RISKS?

The pill is safer than pregnancy. It should not be used within 6 weeks of delivery. After six weeks, it should not impact breastfeeding. Serious problems such as blood clots, are rare. Seek medical attention if you develop the following: pain, swelling, and redness in an arm or leg, weakness or numbness on one side of your body, severe headache, vision problems, chest pain, difficulty breathing, or think you may be pregnant.

This method does not protect against sexually transmitted infections (STIs) or HIV. Always use condoms to protect yourself.

WHAT IF I HAD UNPROTECTED SEX?

If you miss 2 or more pills in a row, you could become pregnant. If it has been less than 5 days after you have had unprotected sex, take emergency contraception (EC). EC works better the sooner you take it.



PROGESTIN-ONLY (POP) OR MINI-PILL

91% EFFECTIVE

HOW DOES THE PROGESTIN-ONLY PILL WORK?

The progestin-only pill (POP) contains a hormone (progestin) like one your body makes. This hormone works by making the mucus in your cervix too thick for sperm to pass through, and thins the lining inside the uterus. It sometimes stops your ovaries from releasing eggs.

HOW DO I START THE PROGESTIN-ONLY PILL?

There are 2 ways to start the progestin-only pill (POP). We recommend the Quick Start Method.

Quick Start: Take the POP as soon as you get the pack. If you take the POP within 5 days after the start of your period, you are protected against pregnancy right away. If you take the POP more than 5 days after the start of your period, you should use condoms as a back-up method for the first 2 days. If you have taken ulipristal acetate emergency contraception (EC), wait 5 days before starting the progestin-only pill.

Next Period: Take the pill within 5 days of the start of your next period.

HOW DO I USE THE PROGESTIN-ONLY PILL?

Take 1 POP each day. After you finish a pack of pills, start a new pack the next day. It is important to take the POP at the same time every day. There is a POP that is available over-the-counter and without a prescription.

WHAT IF I MISS A PROGESTIN-ONLY PILL?

I forgot **ONE** pill: Take one pill as soon as you can. Take your next pill at the usual time, even if that means taking 2 pills on the same day. If you take a pill more than 3 hours late, use condoms as a back-up method for the next 2 days.

I forgot **TWO** or more pills: Take one pill as soon as you can. Take your next pill at the usual time, even if that means you take 2 pills on the same day. Use condoms for the next 2 days as a back-up method.

WHAT CAN I EXPECT?

You will feel about the same.

The following are common side effects: You may have more bleeding between periods and menstrual changes compared to the combined progestin and estrogen pill.

DOES THE PROGESTIN-ONLY PILL HAVE BENEFITS?

The POP is safe and effective birth control.

It is safe to use while breastfeeding. The POP has no effect on your ability to get pregnant in the future.

DOES THE PROGESTIN-ONLY PILL HAVE RISKS?

The POP is safer than pregnancy. Serious problems are rare. Seek medical attention if you develop the following: severe headaches, very heavy bleeding, or think you might be pregnant.

This method does not protect against sexually transmitted infections (STIs) or HIV. Always use condoms to protect yourself.

WHAT IF I HAD UNPROTECTED SEX?

If you miss a POP by more than 3 hours, you could become pregnant.

If it has been less than 5 days after you have had unprotected sex, take emergency contraception (EC). EC works better the sooner you take it.

THE RING

91% EFFECTIVE



HOW DOES THE RING WORK?

The ring contains hormones (estrogen and progestin) like the ones your body makes. These hormones stop your ovaries from releasing eggs, make the mucus in your cervix too thick for sperm to pass through, and thin the lining inside the uterus. There is a ring you use for a month, and a reusable ring that you use for a year.

HOW DO I START THE RING?

There are 2 ways to start the ring. We recommend the Quick Start Method for the monthly ring and Next Period for the reusable ring.

Quick Start: Put in the ring as soon as you get the pack. If you put the ring in on the first day of your period, you are protected against pregnancy right away. If you insert the ring from the second day or later of your period, you should use condoms as a back-up method for the first 7 days. If you have taken ulipristal acetate emergency contraception (EC), wait 5 days before starting the ring.

Next Period: Put the ring in on the first day of your next period. Put the annual reusable ring in between the second and fifth days of your period. If your periods are not regular or if you start using the annual reusable ring more than 5 days from when you started your period, you should use a back-up method for the first 7 days.

HOW DO I USE THE RING?

The ring is a small, bendable, plastic circle that you insert into your vagina. There is no perfect way to insert the ring; you just place it into your vagina to where you can no longer feel it. You leave the ring in your vagina for 3 weeks, and remove it for the 4th week. Remove the ring by hooking a finger under the rim and pulling it out. Most people get their period during the ring-free week. Insert a new ring monthly at the end of the 4th week.

You can store the ring at room temperature for up to four months or in the refrigerator, where it lasts longer. When using the annual reusable ring, remove it for the 4th week, and wash it with mild soap and warm water and pat dry with a clean towel. You should store the ring at room temperature. At the end of the ring-free week, wash the ring again before inserting it in your vagina.

WHAT IF I FORGET TO CHANGE THE RING?

If you have left the ring in for more than 3 weeks but less than 4 weeks, remove the ring to have your period. Put a new ring in 7 days later. If you have left the ring in for 4 weeks or longer, put a new ring in immediately, and use a back-up method for the next 7 days.

WHAT IF THE RING COMES OUT?

You should insert it again as soon as possible and leave it in until the scheduled ring removal day. If it has been less than 48 hours, no back-up condoms are needed. If the ring has been out for more than 48 hours, use condoms for the next 7 days.

WHAT CAN I EXPECT?

You will feel about the same. The following are common side effects: In the first few months you may have nausea, bleeding between periods, and/or breast discomfort. These problems generally go away after 2-3 months. Some people are concerned about weight gain. This is uncommon with the ring. Some people report a change in their mood or headaches. These can be better or worse.

DOES THE RING HAVE BENEFITS?

The ring is safe and effective birth control. Periods may be regular, lighter, less painful, and shorter. You may have less acne. The ring has no effect on your ability to get pregnant in the future. You don't have to remember to use it every day.

DOES THE RING HAVE RISKS?

The ring is safer than pregnancy. It should not be used within 6 weeks of delivery; afterwards it should not impact breastfeeding. Serious problems, such as blood clots, are rare. Seek medical attention if you develop the following: pain, swelling, and redness in an arm or leg, weakness or numbness on one side of your body, severe headaches, vision problems, chest pain, difficulty breathing, or think you might be pregnant. This method does not protect against sexually transmitted infections (STIs) or HIV. Always use condoms to protect yourself.

WHAT IF I HAD UNPROTECTED SEX?

If the annual ring (Annovera) is out for more than 2 hours or the monthly ring (Nuvaring/EluRyng) is out for more than 3 hours, or you don't change the ring for 4 weeks or longer, you could become pregnant. If the ring is out for more than 48 hours, or you don't change the ring for 4 weeks or longer, you could become pregnant. If it has been less than 5 days after you have had unprotected sex, take emergency contraception (EC). EC works better the sooner you take it.



THE PATCH

91% EFFECTIVE

HOW DOES THE PATCH WORK?

The patch contains hormones (estrogen and progestin) like the ones your body makes. These hormones stop your ovaries from releasing eggs, make the mucus in your cervix too thick for sperm to pass through, and thin the lining inside the uterus.

HOW DO I START THE PATCH?

There are 2 ways to start the patch. We recommend the Quick Start Method.

Quick Start: Put on the patch as soon as you get the pack. If you put on the patch within 5 days after the start of your period, you are protected against pregnancy right away. If you put on the patch more than 5 days after the start of your period, you should use condoms as a backup method for the first 7 days. If you have taken ulipristal acetate emergency contraception (EC), wait 5 days before starting the patch.

Next Period: Put on the patch within 5 days of the start of your next period.

HOW DO I USE THE PATCH?

The patch is a thin square or circle of sticky plastic that you wear on your skin. You wear each patch for one week. You can put it on your buttocks, lower abdomen, or upper back. Place the patch on a clean, dry area and make sure the edges stick well. Do not use lotions or creams before putting on the patch. You will use a new patch every week for 3 weeks and no patch for the 4th week, when you can expect to have a period. Start a new box of patches at the end of the 4th week.

WHAT IF THE PATCH FALLS OFF?

If the patch comes off, apply a new patch as soon as possible. Change your patch on the same patch change day. If the patch was off for less than 48 hours, no backup condoms are needed. If the patch has been off for more than 48 hours, use condoms for the next 7 days.

WHAT IF I FORGET TO CHANGE THE PATCH?

The patch has enough hormones for 9 days. If you leave the patch on for up to 9 days, put on a new patch. If you leave the patch on for more than 9 days, put on a new patch and use condoms as a backup method for the next 7 days.

WHAT CAN I EXPECT?

You will feel about the same.

The following are common side effects: In the first few months you may have nausea, bleeding between periods, and/or breast discomfort. These problems generally go away after 2-3 months. Some people are concerned about weight gain. This is uncommon with the patch. Some people report a change in their mood or headaches. These can be better or worse.

Some people get skin irritation from the patch. To prevent this, put the patch in a new spot each week.

DOES THE PATCH HAVE BENEFITS?

The patch is safe and effective birth control. Periods may be regular, lighter, less painful, and shorter. You may have less acne. The patch has no effect on your ability to get pregnant in the future. You don't have to remember to use it every day.

DOES THE PATCH HAVE RISKS?

The patch is safer than pregnancy. It should not be used within 6 weeks of delivery, afterwards it should not impact breastfeeding. The patch has more estrogen than most combined birth control pills and may increase your risk of blood clots. However, serious problems, such as blood clots, are rare. Seek medical attention if you develop the following: pain, swelling, and redness in an arm or leg, weakness or numbness on one side of your body, severe headaches, vision problems, chest pain, difficulty breathing, or think you might be pregnant.

This method does not protect against sexually transmitted infections (STIs) or HIV. Always use condoms to protect yourself.

WHAT IF I HAD UNPROTECTED SEX?

If your patch is off for more than 48 hours, or if you leave it on for more than 9 days, you could become pregnant. If it has been less than 5 days after you have had unprotected sex, take emergency contraception (EC). EC works better the sooner you take it.